Risk Factors Among High-Cost Homeless Populations

Data Tool	What it is Used For	How Data Gathered	Risk Areas Identified	Scoring
10 th Decile	Predict 10% Highest-	Questionnaire	Mental Illness Combined w/Incarceration (strongest	.35 or Higher:
(Los Angeles,	Cost Homeless People:	Administered in	predictor, particularly psychosis , personality disorder)	Incurring 10%
Economic	Eligible for "10 th -Decile	Hospitals (27 data		Highest
Roundtable)	Project" Services*	points if	Hospital Inpatient Admission in Last 3 Years	Hospital Costs
		incorporates		
		incarceration; 51	Over 46 Years Old	
		points if includes		
		only health care	Chronic Medical Disease: HTN , Respiratory Disease ,	
		history)	Neurological	
			Override Tool if Homeless Plus: HIV, Chronic Liver Disease	
			& Cirrhosis, Venous Embolism, Drug-Induced Mental D/O,	
			Cardiac Dysrhythmia, Poisoning by Central Nervous Sys	
D. H. H.	A1 21 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		Stimulants, Disease of Pancreas, Pulmonary TB	50 U. I
Predicting	Algorithm to Identify	Uses Hospital	No Usual Source of Care	50 or Higher
Hospital Readmissions	Medicaid Patients at	Administrative Data	FR Visits and Innetions Admissions in Last 2 Vacus	Indicates High
(New York	High Risk of Hospital Readmission in Next 12	Based on Diagnoses, Service Utilization	ER Visits and Inpatient Admissions in Last 3 Years	Risk of Hospital Readmission
University-	Months	History in Prior 3	Homelessness	(with 100 being
Bellevue	IVIOIILIIS	Years	riomeiessness	highest score)
Hospital Pilot)		Tears	Conditions: Mental Illness, Substance Use, HTN,	riigriest score)
1103pitai i iiotj			COPD/Asthma, Seizure D/O, Cirrhosis, Coronary Artery	
			Disease, Deep Vein Thrombosis/Embolus	
Coordinated	Composite Database of	Integration of	For Top 1% of Health Care Utilizers, Risk Factors Include:	Top 1%
Case	Integrated Medical,	Multiple City-County	Homelessness (17.8% of total cohort, 69.5% of top 1%)	
Management	Psychological, and	Databases		
System (San	Social Info About High-		Elixhauser Conditions (conditions predicting early	
Francisco	Risk, Complex,		mortality) with Highest Frequency Among Top 1%: Cardiac	
Dept. of Public	Vulnerable Populations		Arrythmias, COPD, DM, Neurological Conditions,	
Health)	Served by San		Electrolyte D/O, HTN, Liver Disease, Alcohol/Drug Abuse,	
	Francisco Dept. of		Depression, Psychoses .	

	Public Health		Combination of These Conditions	
PRISM (Predictive Risk Intelligence SysteM) Washington State	Chronic Care Management Support Tool: Eligibility for Care Management Interventions for Dual- Eligible Population	Claims Data, CARE (functional) Assessment	Diagnosis by Severity, by Combination Pharmacy Data High Admission Rates to Hospitals	1.5 or Higher: Expected to Have Medicaid Costs That Are 50% Higher than Average Over Next 12 Months (on a PMPM basis)
Common Risk Factors for High-Costs:			Homelessness Admission to Hospitals Over Last 3 Years Combination of Conditions Common Conditions: • Mental Illness (esp. psychoses) • Substance Use • HTN • Cardiovascular Conditions (Dysrrythmias) • Respiratory Conditions Liver Disease	

^{*}The 10th-Decile Project Tool specifically seeks to exclude people who need nursing home care, since it is intended to determine eligibility for services and housing for people who can live independently. People are screened out if determined risk of need for nursing home care based on these criteria: need assistance in moving in and out of wheelchair, colostomy bag, urinary catheter, tracheotomy, intravenous therapy, serious wounds that require ongoing wound care.