

## Topical Antimicrobial Agent Skin Reaction Report

Patient ID: \_\_\_\_\_

Birthweight: \_\_\_\_\_ grams EGA: \_\_\_\_\_ weeks Age at Event: \_\_\_\_\_ days

### Antimicrobial Agent(s) Used:

- Chlorhexidine Preparations
  - Alcohol:  Chloraprep  Chlorascrub
  - Aqueous:  2% Concentration (Exocrine & other brands)  4% Preparation
  - Other concentration: \_\_\_\_\_ (Specify name)
- Povidone Iodine (Betadine<sup>®</sup>, etc)
- Alcohol
- Other agent: Specify \_\_\_\_\_

### Style of applicator:

- Prep pad
- SEPP (squeeze & pop vial-like applicator)
- Sponge
- Swab stick
- Gauze

### Agent Removal:

- Not removed
- Removed after procedure
- Unknown

### Removal Agent:

- Water
- Saline
- Alcohol

### Method of Removal:

- Not removed
- Saturated prep pad
- Gauze

### Area Where Agent Applied & Reaction Noted:

- Umbilical area
  - Did agent pool after application?
    - On trunk  On back
- Extremity  Arm  Leg
- Heel
- Scalp
- Other: \_\_\_\_\_

### Procedure Performed:

- Vascular catheter insertion
  - PICC
  - Umbilical
  - PIV
  - Peripheral artery line
- LP
- Other: Specify \_\_\_\_\_

### Time Reaction Noted:

- Upon application
- After application: \_\_\_\_\_ Minutes \_\_\_\_\_ Hours \_\_\_\_\_ Days

### Description of Skin Reaction (check all that apply):

- Erythema
- Edema
- Drying
- Flaking
- Cracking
- Excoriation
- Weeping
- Raised, rash-like
- Other: Describe \_\_\_\_\_
- Could reaction be due to dressing applied over catheter?
  - Yes  No  Unsure

### Treatment Provided:

- None
- Kept dry
- Emollient/antimicrobial agent applied topically
- Wound care product used: Specify \_\_\_\_\_
- Other: Specify \_\_\_\_\_